arXiv:1204.2498v1 [math.OC] 11 Apr 2012

A closed form solution for a stochastic control problem with quasi-polynomial value function*

Peter Kratz†

March 2, 2013

Abstract

We study a constrained stochastic control problem with jumps; the jump times of the controlled process are given by a Poisson process. The cost functional comprises quadratic components for an absolutely continuous control and the controlled process and an absolute value component for the control of the jump size of the controlled process. We characterize the value function by a quasi-polynomial of degree two, i.e., a "polynomial" whose coefficients depend on the state of the system. This ansatz allows us to derive the solution of the optimization problem via dynamic programming and a novel interpolation method in closed form; the value function is a classical solution of the HJB equation corresponding to the problem. The state space is separated by a time dependent boundary into a continuation region where the optimal jump size of the controlled process is positive and a stopping region where it is zero.

1 Introduction

For most non-linear-quadratic control problems, closed form solutions are rather difficult to obtain as the form of the value function cannot be guessed easily. In this paper, we propose an interpolation method for a non-linear-quadratic stochastic control problem whose cost functional comprises both quadratic components and an absolute value component; we derive the solution of the problem in closed form and show that the value function is a classical solution of the corresponding Hamilton Jacobi Bellman (HJB) equation.

Given a control $u = (\xi, \eta)$, the controlled process X^u satisfies the stochastic differential equation

$$dX^u(s) = -\xi(s)ds - \eta(s)d\pi(s), \quad X^u(t) = x$$

and the terminal constraint $\lim_{s\to T-} X^u(s) = 0$; here, $x \in \mathbb{R}$ is the initial state of the system at time $t \in [0, T]$ and π is a Poisson process with intensity $\theta \geq 0$. The costs of u are given by

$$J(t, x, u) := \mathbb{E}\Big[\int_t^T \Lambda\xi(s)^2 + \theta\Gamma|\eta(s)| + \alpha\sigma^2 X^u(s)^2 ds\Big]^1$$

and have to be minimized.

Using dynamic programming, it can be shown that the value function of the corresponding discrete time minimization problem is piecewise a quadratic polynomial (see Kratz and Schöneborn [2012a]): At the last control time $t_N = T$, the value function is quadratic; in order to obtain the optimal control and the value function at time t_{N-1}, the term $|\eta|$ necessitates the distinction of several (more precisely two) cases; the number of cases increases with the number of backward time steps. It is hence natural to characterize the value function of the continuous time analogue by a *quasi-polynomial*: a "polynomial" whose coefficients depend on the state of the system. Plugging this ansatz into the HJB equation allows us to drive a candidate for the value function and the optimal control via extensive heuristic considerations. It turns out that the value function is a true polynomial for small states $|x| \leq \beta(t)$ and for large states $|x| \geq \gamma(t)$ (for functions $0 < \beta \leq \gamma$). For intermediate

---

*I wish to thank Peter Bank, Ulrich Horst, Werner Kratz and Torsten Schöneborn for useful discussions and comments. I am also grateful to seminar participants at Humboldt University Berlin and at German Probability and Statistics Days Mainz. This research was supported by Deutsche Bank through the Quantitative Products Laboratory.

†Humboldt Universität zu Berlin, Unter den Linden 6, 10099 Berlin, Germany. Email: kratz@mathematik.hu-berlin.de

[1] $\Lambda > 0$ and $\Gamma, \alpha, \sigma \geq 0$ are constants motivated by the application of the control problem outlined in Section 6.3.

states ($\beta(t) < |x| < \gamma(t)$) the value function is given by an "interpolation" of these polynomials; we derive an implicit representation of the interpolation function as the inverse of an explicitly known injective function. The interpolation method we use appears promising for the solution of related control problems with both quadratic and absolute value cost components.[2] In the present case, the fact that the candidate of the value function and the optimal control is given in closed form is essential for the verification theorem as it enables us to resolve problems arising due to the singularity of the value function at time T which stems from the terminal constraint.

The optimal control of the jump size of the controlled process, η^*, divides the state space into two regions: a "continuation region" ($|x| > \beta(t)$) where $|\eta^*| > 0$ and a "stopping region" ($|x| \leq \beta(t)$) where $\eta^* = 0$. The structure of the optimal control hence resembles the solution of free boundary problems arising in optimal stopping (see, e.g., Pham [2009]) and in singular control problems connected to optimal stopping (see, e.g., Benth and Reikvam [2004]). The control problem studied in this paper is obviously not a singular control problem. Nevertheless, the similarity of the structure is not surprising due to the "linearity" of the costs for the jump size control η. Similarly, we prove a "smooth-fit" principle for the value function on the boundary β; it is essential in the heuristic derivation of the solution of the optimization problem.

The paper contributes to the literature on stochastic control with jumps; standard references are the book by Øksendal and Sulem [2007] and Hanson [2007]. Several papers study stochastic control problems with similar terminal constraints. Schied et al. [2010] study multi-dimensional CARA utility maximization without jumps. The multi-dimensional generalization of the present problem was solved by Kratz and Schöneborn [2012b] for $\Gamma = 0$; in this case, the optimization problem is linear-quadratic and the difficulty stems from the combination of the terminal constraint and the multi-dimensionality. Finally, Naujokat and Westray [2011] and Naujokat and Horst [2011] solve similar control problems by using the stochastic maximum principle and obtain presentations of the optimal controls via FBSDEs. I am not aware of any other work where the value function is represented by a quasi-polynomial and thus a closed form solution for a non-linear-quadratic stochastic control problem is obtained.

The remainder of the article is structured as follows. In Section 2, we describe the optimization problem rigorously. In Section 3, we derive candidates for the value function and the optimal control in closed form by extensive heuristic considerations. We analyze the candidate value function in Section 4: it is continuously differentiable and a classical solution of the HJB equation with singular terminal condition corresponding to the optimization problem. In Section 5, we execute a verification argument. We conclude in Section 6 by discussing properties of the optimal control; furthermore, we motivate the optimization problem by an application to the finance.

2 Optimization problem

In the following, we specify the optimization problem. For a fixed time interval $[0, T]$, we consider the stochastic basis $(\Omega, \mathcal{F}, \mathbb{P}, \mathbb{F} = (\mathcal{F}_t)_{t\in[0,T]})$, where $(\mathcal{F}_t)_t$ is the completion of $\big(\sigma\big(\pi(s)|0 \leq s \leq t\big)\big)_t$ for a

$$\text{Poisson process } \pi = (\pi(t))_{t\in[0,T]} \text{ with intensity } \theta \geq 0.$$

Let $t \in [0, T)$ be a given point in time and $x \in \mathbb{R}$. We call a two-dimensional stochastic process

$$(u(s))_{s\in[t,T)} = (\xi(s), \eta(s))_{s\in[t,T)}$$

a *control* if ξ is progressively measurable and η is predictable. Given a control u, the *controlled process* at time $s \in [t, T)$ is given by the solution of the following controlled stochastic differential equation:

$$dX^u(s) = -\xi(s)ds - \eta(s)d\pi(s), \quad X^u(t) = x \tag{1}$$

such that the left hand side in (1) is well-defined. In view of Equation (1) (and somewhat unconventionally), we denote ξ as the *absolutely continuous control* and η the *jump control* (note that the jump size and not the jump time is controlled). We impose the following conditions.

Definition 2.1. Let $t \in [0, T)$ and $x \in \mathbb{R}$ be fixed. Let $u = (u(s))_{s\in[t,T)} = (\xi(s), \eta(s))_{s\in[0,T)}$ be a control, i.e., ξ is progressively measurable and η is predictable.
We call u an *admissible control* (and write $u \in \mathbb{A}(t, x)$) if it fulfills the following conditions.

---

[2] The application of the method to related control problems requires significant adjustments in order to account for problem specific properties, cf. Section 3.

(i) The Stochastic Differential Equation (1) possesses a unique solution on $[t,T)$.

(ii)

$$\mathbb{E}\Big[\int_t^T |\xi(s)|^4 ds\Big] < \infty, \quad \mathbb{E}\Big[\int_t^T |\eta(s)|^8 ds\Big] < \infty.$$

(iii) If $\theta = 0$, then $\eta(s) = 0$ for all $s \in [t,T)$.

(iv) $\lim_{s\to T-} X^u(s) = 0 \quad \text{a.s.}$

Let us shortly comment on Definition 2.1. Condition (ii) is required for the moment bound in Proposition 2.2 which is in turn needed for the verification argument later. Condition (iii) is needed in order to ensure uniqueness of optimal controls (cf. the dynamics of the controlled process in Equation (1) and the Cost Functional (2)). The terminal constraint (iv) is motivated the application of the control problem outlined in Section 6.3; the terminal constraint can be replaced by any other real number without changing the control problem essentially. We readily observe that Definition 2.1 is satisfied if $\xi(s) = \frac{x}{T-t}$ and $\eta(s) = 0$ for all $s \in [t,T)$; hence $\mathbb{A}(t,x) \neq \emptyset$.

Definition 2.1 implies the following moment estimate for the controlled process. We need this result later for the proof of the verification theorem in Section 5 (the assertion can be proved by applying standard arguments, see Kratz and Schöneborn [2012b], Proposition 2.5).

Proposition 2.2. *Let $t \in [0,T)$, $x \in \mathbb{R}$ and $u \in \mathbb{A}(t,x)$. Then*

$$\mathbb{E}\Big[\sup_{t\le s\le T} |X^u(s)|^4\Big] < \infty, \quad \textit{in particular} \quad \mathbb{E}\Big[\int_t^T |X^u(s)|^4 ds\Big] < \infty.$$

We assume the following costs for an admissible strategy u.

$$J(t,x,u) := \mathbb{E}\Big[\int_t^T f(\xi(s),\eta(s),X^u(s))ds\Big], \tag{2}$$

where $f : \mathbb{R}^n \times \mathbb{R}^n \to \mathbb{R}$, $f(\xi,\eta,x) := \Lambda\xi^2 + \theta\Gamma|\eta| + \alpha\sigma^2x^2$ for $\Lambda > 0$ and $\Gamma, \alpha\sigma \geq 0$. The goal of this paper is to solve the following cost minimization problem:

$$v(t,x) := \inf_{u\in\mathbb{A}(t,x)} J(t,x,u). \tag{OPT}$$

We observe that $\xi(s) = \frac{x}{T-t}$, $\eta(s) = 0$ results in finite costs; hence $v(t,x) < \infty$ and the optimization problem is well-defined.

The key step towards the solution of the Optimization Problem (OPT) is the derivation of a candidate $w : [0,T) \times \mathbb{R} \longrightarrow \mathbb{R}_+$ for the value function. Heuristic considerations suggest that w should satisfy the following HJB equation:

$$\begin{aligned} \frac{\partial w}{\partial t}(t,x) &= \sup_{u=(\xi,\eta)\in\mathbb{R}\times\mathbb{R}} \Big[\theta\big(w(t,x) - w(t,x-\eta)\big) + \frac{\partial w}{\partial x}(t,x)\xi - f(\xi,\eta,x)\Big], \\ \lim_{t\to T-} w(t,x) &= \begin{cases} 0 & \text{if } x = 0 \\ \infty & \text{else.} \end{cases} \end{aligned} \tag{HJB}$$

The singularity of the HJB Equation (HJB) is due to the terminal constraint Definition 2.1 (iv); it causes difficulties for the verification argument as we have to consider the limit $s \to T-$. In order to resolve this problem, it is desirable to find a solution of (HJB) in closed form. For the case $\Gamma = 0$, the multi-dimensional analogue of the Optimization Problem (OPT) was solved by Kratz and Schöneborn [2012b] by plugging a quadratic ansatz into (HJB). In this case, a classical solution of the respective HJB equation is given by a quadratic form which is characterized by the principal solution of a matrix differential equation; although an explicit solution for this differential equation is only known for the one-dimensional case, explicit upper and lower bounds for the solution resolve the problems associated with the singular terminal condition. In the case of this paper, a quadratic ansatz fails; the term $\theta\Gamma|\eta|$ renders the optimization problem non-linear-quadratic. In order to find a candidate for the solution of the HJB Equation (HJB), more elaborate considerations are required; this is accomplished heuristically in Section 3 and verified in Section 4.

3 Heuristic derivation of the candidate value function

Let us for now assume that $x > 0$ and $\alpha\sigma^2 > 0$. The results are symmetric and can directly be transferred to the case $x < 0$; the case $\alpha\sigma^2 = 0$ has a somewhat simpler structure and can be obtained by straightforward modifications or by taking the limit $\alpha\sigma^2 \to 0$ (cf. Section 6.2). Kratz and Schöneborn [2012a] solve the discrete time analogue of the Optimization Problem (OPT) in closed form. If $\Gamma > 0$, the value function is not quadratic in x because of the term $|\eta|$; instead it is piecewise a polynomial of degree two. The authors prove this result by a backward induction using dynamic programming. At the last control time t_N, the value function is quadratic. In order to obtain the optimal control and the value function at time t_{N-1}, the term $|\eta|$ necessitates the distinction of several (more precisely two) cases; the number of cases increases with the number of backward time steps: one additional case per step. The number of polynomials which are glued together in order to obtain the value function hence depends on the number of trading times until t_N. In a similar way, the optimal control is piecewise affine linear in x. Intuitively, a similar structure should hold for a larger class of optimization problems with quadratic and absolute value cost components.

The structure of the value function and the optimal control in the discrete time case gives us certain hints about the structure of the solution of the continuous time Optimization Problem (OPT). The continuous time model can be regarded as the limit of the discrete time model as the number of trading times tends to infinity. In this sense, the finitely many quadratic polynomials forming the value function in the discrete time case become infinitely many for the continuous time case; these must be merged together in some way. We establish an interpolation method accomplishing this task: a candidate of the value function is identified in closed form and we call its representation as a "polynomial" with state-dependent coefficients a *quasi-polynomial* in the following. In view of the solution of the discrete time optimization problem, this is the natural representation for the value function.

As a first step, we require the results from Kratz and Schöneborn [2012b] for the case $\Gamma = 0$.[3] In this case, the value function v and the optimal control $u^* = (\xi^*, \eta^*)$ are given by

$$v(t,x;\theta) = C(t;\theta)x^2, \quad \xi^*(t,x;\theta) = \frac{C(t,x,\theta)}{\Lambda}x, \quad \eta^*(t,x;\theta) = \mathbb{1}_{\{\theta>0\}}x, \tag{3}$$

where

$$C(t;\theta) = \frac{\Lambda\tilde{\theta}}{2}\coth\Big(\frac{\tilde{\theta}}{2}(T-t)\Big) - \frac{\Lambda\theta}{2} \quad \text{for} \quad \tilde{\theta} := \sqrt{\theta^2 + \tfrac{4\alpha\sigma^2}{\Lambda}}. \tag{4}$$

We make the following heuristic observations about the structure of the value function and the optimal control which are partly inspired by the discrete time case.

- For small initial states x, the absolute value costs $\theta\Gamma|\eta|$ outweigh the quadratic costs $\Lambda\xi^2$. Hence, η should be zero and the value function should satisfy $v(t,x) = C(t;0)x^2$ (cf. Equation (4)). Also the optimal control should be given as in (3) for $\theta = 0$.
- For large initial states x, the value function is a quadratic polynomial in the discrete time case (as there are only finitely many polynomials forming the value function). As the number of time steps approaches infinity, this property could in principle break down. The insights from the discrete time optimization problem suggest however that this is not the case; we hence make the educated guess that the value function is a quadratic polynomial and that the optimal control is affine linear in x.
- For intermediate states, we "interpolate" the value function in such a way that it is continuously differentiable.

In Section 3.1, we make the ansatz that the value function is a quasi-polynomial of degree two. In Section 3.2, we deduce a candidate for the optimal control (dependent on the coefficients of this quasi-polynomial) using the HJB Equation (HJB) and obtain a candidate for the boundary below which $\eta = 0$ is optimal. In Section 3.3, we derive differential equations for the coefficients of the candidate value function.

3.1 Quasi-polynomial ansatz for the candidate value function

We make the ansatz that a candidate w for the value function is a quasi-polynomial of degree two:

$$w(t,x) = \bar{C}_1(t,x)x^2 + \bar{C}_2(t,x)|x| + \bar{C}_3(t,x) \tag{5}$$

---

[3] The important special case $\theta = 0$ was first solved by Almgren and Chriss [2001] and later by Schied et al. [2010].

for functions $\bar{C}_i : [0,T) \times \mathbb{R} \to \mathbb{R}$, $i = 1,2,3$ (cf. the discussion above).

In the following we reflect on the optimal control. These considerations yield possible properties of the coefficients of w as in Equation (5). Assume that $u^*(t,x) = (\xi^*(t,x), \eta^*(t,x))$ is the optimal control. For small states, the absolute value costs $\theta\Gamma|\eta|$ are larger than the quadratic costs $\Lambda\xi^2$. Therefore, there should be a time dependent boundary $\beta : [0,T) \to \mathbb{R}_+$ such that

$$\eta^*(t,x) = 0 \quad \text{for } |x| \le \beta(t). \tag{6}$$

For larger states, the absolute value costs are smaller than the quadratic costs; therefore we expect that the control η is such that a jump of π decreases the controlled process X^u to the level β, where further controls $\eta > 0$ are too costly:

$$\eta^*(t,x) = \operatorname{sgn}(x)\big(|x| - \beta(t)\big) \quad \text{for } |x| > \beta(t). \tag{7}$$

As $\eta^* = 0$ for $|x| \le \beta(t)$, the optimal absolutely continuous control below the boundary should be the one given in Equation (3) for $\theta = 0$, i.e, the value function and the optimal control are

$$w(t,x) = C(t;0)x^2, \quad \xi^*(t,x) = \frac{C(t;0)}{\Lambda}x, \quad \eta^*(t,x) = 0, \tag{8}$$

respectively. In other words, we expect the coefficients of w as in Equation (5) to fulfill $\bar{C}_1(t,x) = C(t;0)$, $\bar{C}_2(t,x) = \bar{C}_3(t,x) = 0$ for $|x| \le \beta(t)$. We illustrate these heuristics in the left and the middle picture of Figure 1.

Let us now assume that these considerations are true. We hope that the value function is differentiable on $[0,T) \times \mathbb{R}$. We use this property for the proof of the verification theorem via the HJB Equation (HJB) (cf. Theorem 5.4 below). A necessary condition for continuity of w at $x = \beta(t)$ is $\bar{C}_1(t,\beta(t)) = C(t;0)$, $\bar{C}_2(t,\beta(t)) = \bar{C}_3(t,\beta(t)) = 0$. In order to be differentiable at $x = \beta(t)$, the right-hand and the left-hand partial derivative with respect to x must be equal, i.e., for $t \in [0,T)$, $x = \beta(t)$,

$$2C(t;0)x = 2\bar{C}_1(t,x)x + \bar{C}_2(t,x) + x^2\frac{\partial \bar{C}_1}{\partial x}(t,x) + x\frac{\partial \bar{C}_2}{\partial x}(t,x) + \frac{\partial \bar{C}_3}{\partial x}(t,x),$$

and therefore a necessary condition for differentiability is

$$\beta(t)^2\frac{\partial \bar{C}_1}{\partial x}(t,\beta(t)) + \beta(t)\frac{\partial \bar{C}_2}{\partial x}(t,\beta(t)) + \frac{\partial \bar{C}_3}{\partial x}(t,\beta(t)) = 0.$$

We make the educated guess that this condition also holds for $x \ne \beta(t)$ (note that it certainly holds for small $x < \beta(t)$ and large x as in these cases the coefficients are constant in x):

$$x^2\frac{\partial \bar{C}_1}{\partial x}(t,x) + |x|\frac{\partial \bar{C}_2}{\partial x}(t,x) + \frac{\partial \bar{C}_3}{\partial x}(t,x) = 0 \tag{9}$$

for all $t \in [0,T)$, $x \in \mathbb{R}$ and thus

$$\frac{\partial w}{\partial x}(t,x) = 2\bar{C}_1(t,x)x + \operatorname{sgn}(x)\bar{C}_2(t,x). \tag{10}$$

3.2 The candidate optimal absolutely continuous control

We assume from now on that the value function is given by w as in Equation (5) respectively as in Equation (8), that all partial derivatives exist, that $\frac{\partial w}{\partial x}$ is given as in Equation (10) and that the optimal control η^* is given by Equations (6) and (7).

We consider the HJB Equation (HJB) and maximize the function

$$\begin{aligned} h(t,x,\xi,\eta) &:= \theta\big(w(t,x) - w(t,x-\eta)\big) + \frac{\partial w}{\partial x}(t,x)\xi - \Lambda\xi^2 - \theta\Gamma|\eta| - \alpha\sigma^2x^2 \\ &= \theta w(t,x) - \theta\big(w(t,x-\eta) + \Gamma|\eta|\big) - \alpha\sigma^2x^2 - \Lambda\Big(\frac{\frac{\partial w}{\partial x}(t,x)}{2\Lambda} - \xi\Big)^2 + \frac{\frac{\partial w}{\partial x}(t,x)^2}{4\Lambda} \end{aligned}$$



Figure 1: Illustrations of the heuristics about the optimal control. In all pictures, the (lower) dashed line denotes the time dependent boundary $\beta(t)$ below which $\eta^* = 0$. Solid lines denote realized trajectories of the controlled process. The left picture shows the scenario where the initial state is below β. The middle picture shows a state which is initially above β. In the displayed scenario, the Poisson process jumps at time τ. The third picture shows the optimal control for two states when the Poisson process never jumps. For the smaller initial state, the controlled process crosses the boundary β at time s, i.e., for $t \leq s$ the smaller solid line represents the function $\bar{X}(t,s)$. The larger trajectory does not cross β during the entire horizon; it is above the function $\bar{X}(t,T)$ which is displayed by the second dashed line.

in (ξ,η). Using Equation (10), we obtain that h is maximal for

$$\xi^* := \xi^*(t,x) := \frac{2\bar{C}_1(t,x)x + \operatorname{sgn}(x)\bar{C}_2(t,x)}{2\Lambda} \tag{11}$$

and for $\eta^* = \eta^*(t,x)$ such that $\bar{h}(\eta) := w(t,x-\eta) + \Gamma|\eta|$ is minimal. For $x > \beta(t)$, Equation (7) suggests that this should be the case for

$$\eta^* = \eta^*(t,x) = x - \beta(t). \tag{12}$$

Furthermore for $\eta^* > 0$, $\bar{h}'(\eta^*) = -\frac{\partial w}{\partial x}(t,x-\eta^*) + \Gamma = 0$ if and only if $\frac{\partial w}{\partial x}(t,x-\eta^*) = \Gamma$. Combining these observations, we obtain the following candidate for the boundary $\beta(t)$:

$$\beta(t) = \frac{\Gamma}{2C(t;0)}. \tag{13}$$

3.3 Differential equations for the coefficients

For now, we consider the case $x > \beta(t)$. Given the assumption that w as in Equation (5) solves the HJB Equation (HJB) with maximizer $u^* = (\xi^*,\eta^*)$ for ξ^* and η^* as in Equations (11) and (12), respectively, we obtain

$$\frac{\partial w}{\partial t}(t,x) = h(t,x,\xi^*,\eta^*).$$

Provided that Equations (9) and (13) hold, this implies

$$\begin{aligned}&\frac{\partial \bar{C}_1}{\partial t}(t,x)x^2 + \frac{\partial \bar{C}_2}{\partial t}(t,x)x + \frac{\partial \bar{C}_3}{\partial t}(t,x)\\ &\quad= \Big(\frac{\bar{C}_1(t,x)^2}{\Lambda} + \theta\bar{C}_1(t,x) - \alpha\sigma^2\Big)x^2 + \Big(\bar{C}_2(t,x)\Big(\frac{\bar{C}_1(t,x)}{\Lambda} + \theta\Big) - \theta\Gamma\Big)x + \theta\bar{C}_3(t,x) + \frac{\theta\Gamma^2}{4C(t;0)} + \frac{\bar{C}_2(t,x)^2}{4\Lambda}.\end{aligned}$$

We hence expect that for $|x| > \beta(t)$, the coefficients of the candidate value function fulfill the ordinary differential equations

$$\frac{\partial \bar{C}_1}{\partial t}(t,x) = \frac{\bar{C}_1(t,x)^2}{\Lambda} + \theta\bar{C}_1(t,x) - \alpha\sigma^2, \tag{14}$$

$$\frac{\partial \bar{C}_2}{\partial t}(t,x) = \bar{C}_2(t,x)\Big(\frac{\bar{C}_1(t,x)}{\Lambda} + \theta\Big) - \theta\Gamma, \tag{15}$$

$$\frac{\partial \bar{C}_3}{\partial t}(t,x) = \theta\bar{C}_3(t,x) + \frac{\theta\Gamma^2}{4C(t;0)} + \frac{\bar{C}_2(t,x)^2}{4\Lambda} \tag{16}$$

and that for $x = \beta(t)$,

$$\bar{C}_1(t,x) = C(t;0), \quad \bar{C}_2(t,x) = \bar{C}_3(t,x) = 0. \tag{17}$$

In order to transfer the Differential Equations (14), (15), (16) and the Conditions (17) into initial value problems, we consider the trajectory resulting from the candidate optimal absolutely continuous control ξ^*. Let $t \in [0,T)$, $x \geq \beta(t)$ and let $s \in [t,T)$ be the first time such that the controlled process crosses the boundary $\beta(s)$, provided that the Poisson process does not jump in $[t,T)$. We define

$$\bar{X}(t,s) := x;$$

in other words, the *deterministic* function $\bar{X}(\cdot,s)$ defined on $[t,s]$ is the candidate for the controlled process corresponding to the candidate optimal absolutely continuous control ξ^*, provided the Poisson process never jumps. Note also that this implies

$$\bar{X}(s,s) = \beta(s) = \frac{\Gamma}{2C(s;0)}$$

(cf. Equation (13)). We illustrate the functions $\bar{X}$ in the third picture of Figure 1.

We modify the notation for the coefficients of the candidate value function in the following way. If $t \in [0,T)$, $x \geq \beta(t)$ and if there exists a smallest $s \in [t,T)$ such that the trading trajectory resulting from the candidate optimal trading intensity ξ^* crosses the boundary β at time s without any jump of π, we write

$$C_1(t,s) := \bar{C}_1(t,x), \quad C_2(t,s) := \bar{C}_2(t,x), \quad C_3(t,s) := \bar{C}_3(t,x).$$

Thus, we expect that the new coefficients C_1, C_2 and C_3 solve the following initial value problems

$$\frac{\partial C_1}{\partial t}(\cdot,s) = \frac{C_1(\cdot,s)^2}{\Lambda} + \theta C_1(\cdot,s) - \alpha\sigma^2, \quad C_1(s,s) = C(s;0), \tag{18}$$

$$\frac{\partial C_2}{\partial t}(\cdot,s) = C_2(\cdot,s)\Big(\frac{C_1(\cdot,s)}{\Lambda} + \theta\Big) - \theta\Gamma, \quad C_2(s,s) = 0, \tag{19}$$

$$\frac{\partial C_3}{\partial t}(\cdot,s) = \theta C_3(\cdot,s) + \frac{\theta\Gamma^2}{4C(\cdot;0)} + \frac{C_2(\cdot,s)^2}{4\Lambda}, \quad C_3(s,s) = 0 \tag{20}$$

and that $\bar{X}$ solves the initial value problem (cf. Equation (11))

$$\frac{\partial \bar{X}}{\partial t}(\cdot,s) = -\frac{C_1(\cdot,s)}{\Lambda}\bar{X}(\cdot,s) - \frac{C_2(\cdot,s)}{2\Lambda} = -\xi^*(\cdot,\bar{X}(\cdot,s)), \quad \bar{X}(s,s) = \frac{\Gamma}{2C(s;0)}. \tag{21}$$

4 The candidate value function

Following the heuristic considerations of Section 3, we define the candidate value function w by

$$w(t,x) := C_1(t,g(t,x))x^2 + C_2(t,g(t,x))|x| + C_3(t,g(t,x)), \tag{22}$$

where C_1, C_2 and C_3 are the solutions of the Initial Value Problems (18), (19) and (20), respectively. The function $g : [0,T] \times \mathbb{R} \to [0,T]$ is given explicitly for $|x| \in [0, \frac{\Gamma}{2C(t;0)}] \cup [\bar{X}(t,T), \infty)$ by

$$g(t,x) := \begin{cases} t & \text{if } |x| \in [0, \frac{\Gamma}{2C(t;0)}] \\ T & \text{if } |x| \in [\bar{X}(t,T), \infty), \end{cases} \tag{23}$$

and implicitly for $|x| \in (\frac{\Gamma}{2C(t;0)}, \bar{X}(t,T))$ by

$$\bar{X}(t,g(t,x)) = |x|, \tag{24}$$

where $\bar{X}$ is the solution of the Initial Value Problem (21), i.e., $g(t,\cdot)$ is the inverse function of $\bar{X}(t,\cdot)$. It is not immediately clear that such an inverse function exists and that the function w defined by Equation (22) is well-defined.

The remainder of the section is structured as follows. In Section 4.1, we compute the solutions of the Initial Value Problems (18), (19), (20) and (21) in closed form. This enables us to prove that w is well-defined. In Section 4.2, we prove that w is continuously differentiable and strictly convex in x. Finally, we show that w is a classical solution of the HJB Equation (HJB) in Section 4.3.

In Sections 4 - 6.1, we assume that $\alpha\sigma^2 > 0$. We treat the case $\alpha\sigma^2 = 0$ in Section 6.2.

4.1 Closed form solutions for the coefficients and the trading trajectory

In order that the value function is well-defined, the function g must be well-defined. Therefore, we require that $\bar{X}$ is strictly monotone in s (cf. Equation (24)). We prove this by directly computing the partial derivative of $\bar{X}$ with respect to s and show that it is strictly positive.

We start by giving closed form solutions for the value function coefficients C_1, C_2, C_3 and for $\bar{X}$, which follow from the Initial Value Problems (18), (19), (20) and (21), respectively. Proposition 4.1 treats the case $s \in (t,T)$, the case $s = T$ is treated separately afterwards.

Proposition 4.1. *Let $t \in [0,T)$, $s \in (t,T)$ and define*

$$\mu(s) := \frac{2C(s;0) + \theta\Lambda}{\tilde{\theta}\Lambda}, \quad \kappa(s) := \operatorname{arcoth}(\mu(s)); \tag{25}$$

recall that $\tilde{\theta} = \sqrt{\theta^2 + \frac{4\alpha\sigma^2}{\Lambda}}$. Then

$$C_1(t,s) = \frac{\Lambda\tilde{\theta}}{2}\coth\big(\frac{\tilde{\theta}}{2}(s-t) + \kappa(s)\big) - \frac{\Lambda\theta}{2} > 0, \tag{26}$$

$$C_2(t,s) = \frac{\theta\Gamma}{2\alpha\sigma^2}\Big(-\theta\Lambda + \frac{\Lambda\tilde{\theta}\big(\sinh\big(\frac{\tilde{\theta}}{2}(s-t)\big) + \mu(s)\cosh\big(\frac{\tilde{\theta}}{2}(s-t)\big)\big) - 2C(s;0)\exp\big(\frac{\theta}{2}(t-s)\big)}{\mu(s)\sinh\big(\frac{\tilde{\theta}}{2}(s-t)\big) + \cosh\big(\frac{\tilde{\theta}}{2}(s-t)\big)}\Big) > 0, \tag{27}$$

$$C_3(t,s) = -\int_t^s \exp\big(\theta(t-u)\big)\Big(\frac{\theta\Gamma^2}{4C(u;0)} + \frac{C_2(u,s)^2}{4\Lambda}\Big)du < 0, \tag{28}$$

$$\begin{aligned}\bar{X}(t,s) = {}& \Big(\frac{\Gamma\mu(s)}{2C(s;0)} + \frac{\theta^2\Gamma}{2\tilde{\theta}\alpha\sigma^2}\Big)\sinh\big(\frac{\tilde{\theta}}{2}(s-t)\big)\exp\big(\frac{\theta}{2}(t-s)\big)\\ &+ \Big(\frac{\Gamma}{2C(s;0)} + \frac{\theta\Gamma}{2\alpha\sigma^2}\Big)\cosh\big(\frac{\tilde{\theta}}{2}(s-t)\big)\exp\big(\frac{\theta}{2}(t-s)\big) - \frac{\theta\Gamma}{2\alpha\sigma^2}.\end{aligned} \tag{29}$$

Proof. Note first that for $s \in (t,T)$, $C(s;0) > \sqrt{\alpha\sigma^2\Lambda}$ by Equation (4). Therefore,

$$\Lambda\tilde{\theta} = \sqrt{\Lambda^2\theta^2 + 4\alpha\sigma^2\Lambda} < \Lambda\theta + 2\sqrt{\alpha\sigma^2\Lambda} \le \Lambda\theta + 2C(s;0);$$

thus $\mu(s) > 1$ and $\kappa(s)$ as in Equation (25) is well-defined for all $s \in [0,T)$.

Equation (26) follows directly as (18) is an initial value problem for a Riccati equation with constant coefficients. (19), (20) and (21) are initial value problems for inhomogeneous linear differential equations whose solutions are readily computed (note that $\frac{1}{C(\cdot;0)}$ is continuous). In order to obtain Equations (27), (28) and (29), elementary but tedious algebraic manipulations are required which can be executed by computer algebra software.[4] □

As $C(T;0)$ is not defined and $\lim_{s\to T-} C(s;0) = \infty$, $C_1(t,s)$, $C_2(t,s)$, $C_3(t,s)$ and $\bar{X}(t,s)$ ($t \in [0,T)$) as in Equations (26) - (29) are not defined for $s = T$. However, we require these objects for the definition of the candidate value function w. We define $C_i(t,T) := \lim_{s\to T-} C_i(t,s)$ and $\bar{X}(t,T) := \lim_{s\to T-} \bar{X}(t,s)$. Elementary considerations verify that these limits exist and are given by

$$C_1(t,T) = \frac{\Lambda\tilde{\theta}}{2}\coth\big(\frac{\tilde{\theta}}{2}(T-t)\big) - \frac{\Lambda\theta}{2} > 0, \tag{30}$$

$$C_2(t,T) = \frac{\theta\Gamma\Lambda}{2\alpha\sigma^2}\Big(\tilde{\theta}\coth\big(\frac{\tilde{\theta}}{2}(T-t)\big) - \tilde{\theta}\frac{\exp\big(\frac{\theta}{2}(t-T)\big)}{\sinh\big(\frac{\tilde{\theta}}{2}(T-t)\big)} - \theta\Big) > 0, \tag{31}$$

$$C_3(t,T) = -\int_t^T \exp\big(\theta(t-u)\big)\Big(\frac{\theta\Gamma^2}{4C(u;0)} + \frac{C_2(u,T)^2}{4\Lambda}\Big)du < 0, \tag{32}$$

$$\bar{X}(t,T) = \Big(\frac{\Gamma}{\tilde{\theta}\Lambda} + \frac{\theta^2\Gamma}{2\tilde{\theta}\alpha\sigma^2}\Big)\sinh\big(\frac{\tilde{\theta}}{2}(T-t)\big)\exp\big(\frac{\theta}{2}(t-T)\big) + \frac{\theta\Gamma}{2\alpha\sigma^2}\cosh\big(\frac{\tilde{\theta}}{2}(T-t)\big)\exp\big(\frac{\theta}{2}(t-T)\big) - \frac{\theta\Gamma}{2\alpha\sigma^2}. \tag{33}$$

It follows immediately from the Formulae (26) - (29) that both the coefficients of the candidate value function and the trajectories $\bar{X}$ are continuously differentiable, even analytic.

---

[4] Alternatively, detailed proofs for these equations can be found in Kratz [2011]; this applies also for later equations whose derivation require tedious algebraic manipulations.

Corollary 4.2. *Let $t \in [0,T)$, $s \in (t,T)$. Then C_1, C_2, C_3 and $\bar{X}$ have continuous partial derivatives of any order in (t,s).*

In order to show that the function g given in Equations (23) and (24) and thus the candidate value function is well-defined, we need monotonicity of the function $\bar{X}(t,\cdot)$. Given Equation (29), we can directly compute the partial derivative of $\bar{X}$ with respect to the second argument. Algebraic manipulations confirm that $\bar{X}(t,\cdot)$ is strictly increasing in (t,T) as required (cf. Footnote 4). We can hence prove the central result of this section.

Proposition 4.3. *The candidate value function w given by Equation (22) is well-defined.*

Proof. By the discussion above, $\bar{X}(t,\cdot)$ possesses an inverse on (t,T). Thus, g as in Equations (23) and (24) is well-defined on $[0,T)\times\mathbb{R}$ (cf. also Equation (33)). Furthermore, $g(t,x) = g(t,|x|) \in [t,T]$ for all $t \in [0,T)$, $x \in \mathbb{R}$, and the functions C_i $(i = 1,2,3)$ are well-defined for $t \in [0,T)$, $s \in [t,T]$. □

4.2 Differentiation and convexity of the candidate value function

The goal of this section is to show that the candidate value function w is continuously differentiable. We are also able to compute the partial derivatives $\frac{\partial w}{\partial t}$, $\frac{\partial w}{\partial x}$ and $\frac{\partial^2 w}{\partial x^2}$. It turns out that $\frac{\partial w}{\partial x}$ has indeed the form we assumed in the heuristics in Section 3.1 (cf. Equation (10)). We also show that $\frac{\partial^2 w}{\partial x^2} > 0$ and hence that w is strictly convex in x. We first state the main theorem.

Theorem 4.4. *The candidate value function w given by Equation (22) is continuously differentiable on $[0,T)\times\mathbb{R}$. Furthermore, we have*

$$\frac{\partial w}{\partial x}(t,x) = 2C_1(t,g(t,x))x + \operatorname{sgn}(x)C_2(t,g(t,x)), \tag{34}$$

$$\frac{\partial w}{\partial t}(t,x) = \begin{cases} C'(t;0)x^2 & \text{if } x = \frac{\Gamma}{2C(t;0)} \\ \frac{\partial C_1}{\partial t}(t,g(t,x))x^2 + \frac{\partial C_2}{\partial t}(t,g(t,x))|x| + \frac{\partial C_3}{\partial t}(t,g(t,x)) & \text{else.} \end{cases} \tag{35}$$

Before we proceed with the proof of Theorem 4.4, we show that g is continuously differentiable. We prove this in the following lemma and compute the partial derivatives of g.

Lemma 4.5. *Let $V := \{(t,x) \in \mathbb{R}^2 | t \in (0,T),\ x \in (\bar{X}(t,t), \bar{X}(t,T))\}$. Then g given by Equations (23) and (24) is continuously differentiable on V and*

$$\frac{\partial g}{\partial x}(t,x) = \frac{1}{\frac{\partial \bar{X}}{\partial s}(t,g(t,x))}, \quad \frac{\partial g}{\partial t}(t,x) = -\frac{\frac{\partial \bar{X}}{\partial t}(t,g(t,x))}{\frac{\partial \bar{X}}{\partial s}(t,g(t,x))}. \tag{36}$$

Proof. Consider the open set $U := \{(t,s) \in \mathbb{R}^2 | t \in (0,T), s \in (t,T)\}$ and the continuously differentiable function $h : \mathbb{R}^2 \longrightarrow \mathbb{R}^2$, $h(t,s) := (t, \bar{X}(t,s))^\top$, (cf. Corollary 4.2). Then for all $(t_0,s_0) \in U$, the Jacobian matrix of h is given by

$$Dh(t_0,s_0) := \begin{pmatrix} 1 & \frac{\partial \bar{X}}{\partial t}(t_0,s_0) \\ 0 & \frac{\partial \bar{X}}{\partial s}(t_0,s_0) \end{pmatrix}.$$

Note that $\frac{\partial \bar{X}}{\partial s}(t_0,s_0) \neq 0$ as $\bar{X}(t_0,\cdot)$ is strictly increasing. Therefore $\det Dh(t_0,s_0) = \frac{\partial \bar{X}}{\partial s}(t_0,s_0) \neq 0$, and we can apply the inverse function theorem. Thus, the inverse function h^{-1} of h exists locally and is locally continuously differentiable. However, on $\tilde{V} := h(U) = \{(t,x) \in \mathbb{R}^2 | t \in (0,T),\ x \in (\bar{X}(t,t), \bar{X}(t,T))\}$, we have $h^{-1} = \tilde{g}$ for $\tilde{g}(t,x) := (t, g(t,x))^\top$ globally, and therefore g is continuously differentiable on V.

The first equation in (36) follows from Equation (24) immediately; the second equation follows by differentiating with respect to t on either side of Equation (24). □

It follows directly that w is continuous on $[0,T)\times\mathbb{R}$. We can now prove the main building block of the proof of Theorem 4.4; we show that

$$\frac{\partial C_1}{\partial s}(t,s)\bar{X}(t,s)^2 + \frac{\partial C_2}{\partial s}(t,s)\bar{X}(t,s) + \frac{\partial C_3}{\partial s}(t,s) = 0.$$

This is equivalent to Equation (9) in the heuristics of Section 3.1 and was the key step towards the derivations of the differential equations for the coefficients C_1, C_2 and C_3. Therefore, the construction of the candidate value function relies strongly on this property. Recall that we had good reasons to believe that Equation (9) is true for $|x| \leq \beta(t)$ (or equivalently $s = t$) and for $|x| > \bar{X}(t,T)$ (or equivalently $s = T$). It is somewhat surprising that it is indeed also correct for $t < s < T$.

The second part of the lemma is a similar assertion which is useful for the computation of the second order derivative $\frac{\partial^2 w}{\partial x^2}$ and hence for the proof of the convexity of w below.

Lemma 4.6. *Let $t \in [0,T)$, and $s \in [t,T)$. Then*

(i)

$$a(t,s) := \frac{\partial C_1}{\partial s}(t,s)\bar{X}(t,s)^2 + \frac{\partial C_2}{\partial s}(t,s)\bar{X}(t,s) + \frac{\partial C_3}{\partial s}(t,s) = 0,$$

(ii)

$$b(t,s) := 2\frac{\partial C_1}{\partial s}(t,s)\bar{X}(t,s) + \frac{\partial C_2}{\partial s}(t,s) = 0.$$

Proof. (i) We first show that

$$a(s,s) = \lim_{t \to s-} a(t,s) = 0 \tag{37}$$

by computing the derivatives of C_1, C_2 and C_3 with respect to s. Secondly, we verify that

$$\frac{\partial a}{\partial t}(t,s) = \theta a(t,s).$$

Hence, there exists a constant $C(s)$ ($s \in (0,T)$) such that for all $t \in [0,s)$, $a(t,s) = C(s)\exp(\theta t)$. Equation (37) implies $0 = \lim_{t \to s-} a(t,s) = C(s)\exp(\theta s)$, i.e., $C(s) = 0$, finishing the proof of (i) (cf. Footnote 4 again for details).

(ii) We proceed similarly as in the proof of (i) and show $\lim_{t \to s-} b(t,s) = 0$ and $\frac{\partial b}{\partial t}(t,s) = \left(\frac{C_1(t,s)}{\Lambda} + \theta\right)b(t,s)$, from which the assertion follows.

□

We are now able to prove that w is continuously differentiable.

Proof of Theorem 4.4. Let $t \in [0,T), x \in \mathbb{R}$. We only treat the case $x \geq 0$. The case $x < 0$ follows accordingly. We consider four cases separately.

(i) $x < \frac{\Gamma}{2C(t;0)}$. In this case we have $g(v,y) = v$ for all (v,y) in some neighborhood of (t,x) (cf. Equation (23)). Therefore, $w(v,y) = C(v;0)y^2$ and consequently (recall that $C_1(t,t) = C(t;0)$, $C_2(t,t) = 0$)

$$\frac{\partial w}{\partial x}(t,x) = 2C(t;0)x = 2C_1(t,t)x + C_2(t,t) = 2C_1(t,g(t,x))x + C_2(t,g(t,x)) \tag{38}$$

and

$$\frac{\partial w}{\partial t}(t,x) = C'(t;0)x^2. \tag{39}$$

(ii) $x \in (\frac{\Gamma}{2C(t;0)}, \bar{X}(t,T))$. This is equivalent to $x = \bar{X}(t,s)$, $g(t,x) = s$ for some $s \in (t,T)$. We have $y \in (\frac{\Gamma}{2C(v;0)}, \bar{X}(t,T))$ and therefore $g(v,y) \in (t,T)$ for all (v,y) in some neighborhood of (t,x); thus

$$w(t,y) = C_1(t,g(t,y))y^2 + C_2(t,g(t,y))y + C_3(t,g(t,y)).$$

Using Lemma 4.5 and Lemma 4.6 (i), we obtain

$$\frac{\partial w}{\partial x}(t,x) = 2C_1(t,g(t,x))x + C_2(t,g(t,x)) \tag{40}$$

and

$$\frac{\partial w}{\partial t}(t,x) = \frac{\partial C_1}{\partial t}(t,g(t,x))x^2 + \frac{\partial C_2}{\partial t}(t,g(t,x))x + \frac{\partial C_3}{\partial t}(t,g(t,x)). \tag{41}$$

(iii) $x > \bar{X}(t,T)$. In this case we have $y > \bar{X}(v,T)$ and $g(v,y) = T$ for all (v,y) in some neighborhood of (t,x) (cf. Equation (23) again); thus $w(v,y) = C_1(v,T)y^2 + C_2(v,T)y + C_3(v,T)$. Therefore,

$$\frac{\partial w}{\partial x}(t,x) = 2C_1(t,T)x + C_2(t,T) = 2C_1(t,g(t,x))x + C_2(t,g(t,x)), \tag{42}$$

$$\frac{\partial w}{\partial t}(t,x) = \frac{\partial C_1}{\partial t}(t,T)x^2 + \frac{\partial C_2}{\partial t}(t,T)x + \frac{\partial C_3}{\partial t}(t,T). \tag{43}$$

(iv) $x \in \{\frac{\Gamma}{2C(t;0)}, \bar{X}(t,T)\}$. Recall first that w is continuous. Furthermore, as $g(t,\cdot)$ is continuous in x with values in $[t,T]$ and C_1 and C_2 are continuous in $[t,T]$, $\frac{\partial w}{\partial x}(t,\cdot)$ can be extended continuously to $\mathbb{R}$ (cf. also Equations (38), (40) and (42)). Let now $h_n \neq 0$ $(n \in \mathbb{N})$, $h_n \to 0$. For large enough n, the mean value theorem implies that there exists ξ_n strictly between x and $x - h_n$ such that $\frac{w(t,x)-w(t,x-h_n)}{h_n} = \frac{\partial w}{\partial x}(t,\xi_n)$. As $\lim_{n\to\infty} \xi_n = x$, this implies

$$\frac{\partial w}{\partial x}(t,x) = \lim_{n\to\infty} \frac{w(t,x) - w(t,x-h_n)}{h_n} = \lim_{n\to\infty} \frac{\partial w}{\partial x}(t,\xi_n),$$

i.e., $w(t,\cdot)$ is differentiable in x with derivative

$$\frac{\partial w}{\partial x}(t,x) = \lim_{n\to\infty} \frac{\partial w}{\partial x}(t,\xi_n) = 2C_1(t,g(t,x))x + C_2(t,g(t,x)).$$

For differentiation with respect to t, note that by the differential equations for $C(t;0)$, $C_1(\cdot,s)$, $C_2(\cdot,s)$ and $C_3(\cdot,s)$, we obtain for $x = \frac{\Gamma}{2C(t;0)}$ that

$$\frac{\partial C_1}{\partial t}(t,t)x^2 + \frac{\partial C_2}{\partial t}(t,t)x + \frac{\partial C_3}{\partial t}(t,t) = C'(t;0)x^2.$$

Therefore, $\frac{\partial w}{\partial t}(\cdot,x)$ can be extended continuously to $[0,T)$ by Equations (39), (41) and (43). Using the mean value theorem in a similar fashion as before, we obtain that $w(\cdot,x)$ is differentiable in t with derivative

$$\frac{\partial w}{\partial t}(t,x) = \begin{cases} \frac{\partial C_1}{\partial t}(t,t)x^2 + \frac{\partial C_2}{\partial t}(t,t)x + \frac{\partial C_3}{\partial t}(t,t) = C'(t;0)x^2 & \text{if } x = \frac{\Gamma}{2C(t;0)} \\ \frac{\partial C_1}{\partial t}(t,T)x^2 + \frac{\partial C_2}{\partial t}(t,T)x + \frac{\partial C_3}{\partial t}(t,T) & \text{if } x = \bar{X}(t,T). \end{cases}$$

□

By Theorem 4.4, we obtain first order partial derivatives of the candidate value function w. We can deduce strict convexity of w in x by computing the second order partial derivative $\frac{\partial^2 w}{\partial x^2}$. The key step towards this goal is Lemma 4.6 (ii).

Theorem 4.7. *The candidate value function w given by Equation* (22) *is strictly convex in $x \in \mathbb{R}$.*

Proof. Let $t \in [0,T)$ and $x \in \mathbb{R}$. We consider the case $x \geq 0$. The case $x < 0$ can be proven accordingly. Thus, $x = \bar{X}(t,s)$ for some $s \in [t,T)$ or $x \geq \bar{X}(t,T)$. In the first case Theorem 4.4, Lemma 4.5 and Lemma 4.6 (ii) yield $\frac{\partial^2 w}{\partial x^2}(t,x) = 2C_1(t,g(t,x))$. For $x > \bar{X}(t,T)$, we obtain $\frac{\partial^2 w}{\partial x^2}(t,x) = 2C_1(t,T)$. Hence, $\frac{\partial^2 w}{\partial x^2}(t,x)$ can be extended continuously to $x = \bar{X}(t,T)$ and we obtain

$$\frac{\partial^2 w}{\partial x^2}(t,\bar{X}(t,T)) = 2C_1(t,T)$$

by the same argument as in the proof of Theorem 4.4. Therefore (and by the respective considerations for $x < 0$), the second partial derivative of w with respect to x exists and $\frac{\partial^2 w}{\partial x^2}(t,x) > 0$, hence w is strictly convex in x. □

4.3 Classical solution of the HJB equation

The fact that w is differentiable and the formulae for the partial derivatives finally enable us to prove that w is a classical solution of the HJB Equation (HJB).

Theorem 4.8. *The candidate value function w given by Equation (22) is a classical solution of the HJB Equation (HJB) with unique maximizer $u^* = (\xi^*, \eta^*)$, where*

$$\xi^* = \xi^*(t,x) := \frac{2C_1(t,g(t,x))x + \operatorname{sgn}(x)C_2(t,g(t,x))}{2\Lambda}, \tag{44}$$

$$\eta^* = \eta^*(t,x) := \begin{cases} \operatorname{sgn}(x)\Big(|x| - \frac{\Gamma}{2C(t;0)}\Big) & \text{if } |x| > \frac{\Gamma}{2C(t;0)} \\ 0 & \text{if } |x| \le \frac{\Gamma}{2C(t;0)}. \end{cases} \tag{45}$$

Proof. The theorem follows directly from the subsequent Lemma 4.9 and Theorem 4.4 by using the Differential Equations (18), (19) and (20) for the case $|x| > \frac{\Gamma}{2C(t;0)}$ and the differential equation $C'(\cdot;0) = \frac{C(\cdot;0)^2}{\Lambda} - \alpha\sigma^2$ for the case $|x| \le \frac{\Gamma}{2C(t;0)}$. ☐

Lemma 4.9. *Let $h : [0,T) \times \mathbb{R} \times \mathbb{R} \times \mathbb{R} \to \mathbb{R}$ be given by*

$$h(t,x,\xi,\eta) := \theta\big(w(t,x) - w(t,x-\eta)\big) + \frac{\partial w}{\partial x}(t,x)\xi - f(\xi,\eta,x).$$

For fixed $t \in [0,T)$ and $x \in \mathbb{R}$, $h(t,x,\cdot,\cdot)$ attains its unique maximum at $u^* = (\xi^*,\eta^*)$ for ξ^* and η^* as in Equations (44) and (45), respectively.

Moreover,

$$h(t,x,\xi^*,\eta^*) = \begin{cases} \Big(\frac{C_1(t,g(t,x))^2}{\Lambda} + \theta C_1(t,g(t,x)) - \alpha\sigma^2\Big)x^2 & \\ \quad + \Big(C_2(t,g(t,x))\Big(\frac{C_1(t,g(t,x))}{\Lambda} + \theta\Big) - \theta\Gamma\Big)|x| & \\ \quad + \theta C_3(t,g(t,x)) + \frac{\theta\Gamma^2}{4C(t;0)} + \frac{C_2(t,g(t,x))^2}{4\Lambda} & \text{if } |x| > \frac{\Gamma}{2C(t;0)} \\ \Big(\frac{C(t;0)^2}{\Lambda} - \alpha\sigma^2\Big)x^2 & \text{if } |x| \le \frac{\Gamma}{2C(t;0)}. \end{cases} \tag{46}$$

Proof. Let $t \in [0,T)$ and $x \in \mathbb{R}$. We have

$$h(t,x,\xi,\eta) = \theta w(t,x) - \alpha\sigma^2 x^2 + h_1(t,x,\xi) + h_2(t,x,\eta)$$

for

$$h_1(t,x,\xi) := -\Lambda\xi^2 + \frac{\partial w}{\partial x}(t,x)\xi = -\Lambda\Big(\frac{\frac{\partial w}{\partial x}(t,x)}{2\Lambda} - \xi\Big)^2 + \frac{\frac{\partial w}{\partial x}(t,x)^2}{4\Lambda}, \quad h_2(t,x,\eta) := -\theta w(t,x-\eta) - \theta\Gamma|\eta|.$$

As $\Lambda > 0$, $h_1(t,x,\cdot)$ is strictly concave in ξ. Furthermore, $h_2(t,x,\cdot)$ is strictly concave in η by Theorem 4.7. Thus, $h(t,x,\cdot,\cdot)$ is strictly concave in (ξ,η) and attains its unique global maximum in (ξ^*,η^*) if $h_1(t,x,\cdot)$ attains its unique global maximum in ξ^* and h_2 attains its unique global maximum in η^*.

By Theorem 4.4, h_1 attains its unique global maximum for ξ^* as in Equation (44).

Note now that $h_2 \le 0$. For $x = 0$, $h_2(t,x,\eta^*) = h_2(t,0,0) = 0$, and therefore h_2 attains its unique global maximum in η^* as in Equation (45).

For $\eta \ne 0$, h_2 is differentiable in η. For $|x| > \frac{\Gamma}{2C(t;0)}$, we have

$$\frac{\partial h_2}{\partial \eta}(t,x,\eta^*) = \theta\Big(\frac{w}{\partial x}(t,x-\eta^*) - \operatorname{sgn}(x)\Gamma\Big) = \theta\Big(2C(t;0)\operatorname{sgn}(x)\frac{\Gamma}{2C(t;0)} - \operatorname{sgn}(x)\Gamma\Big) = 0$$

for $\eta^* = \eta^*(t,x)$ as in Equation (45) (note that $\eta^*(t,x) \ne 0$ for $|x| > \frac{\Gamma}{2C(t;0)}$). By strict concavity, we conclude that $h_2(t,x,\cdot)$ attains its unique maximum at η^* as in Equation (45).

Let now $x \in (0, \frac{\Gamma}{2C(t;0)})$. We have

$$\lim_{\eta \to 0-} \frac{\partial h_2}{\partial \eta}(t,x,\eta) = \lim_{\eta \to 0-} 2\theta C(t;0)(x-\eta) + \theta\Gamma = 2\theta C(t;0)x + \theta\Gamma > 0,$$

$$\lim_{\eta\to 0+}\frac{\partial h_2}{\partial \eta}(t,x,\eta) = \lim_{\eta\to 0+} 2\theta C(t;0)(x-\eta) - \theta\Gamma = 2\theta C(t;0)x - \theta\Gamma < 0.$$

Therefore, $\eta^* = 0$ maximizes h_2 uniquely by strict concavity as required.

By continuity of η^* and h_2, $\eta^*(t,x) = 0$ maximizes h_2 uniquely for $x = \frac{\Gamma}{2C(t;0)}$.

The case $x \in [-\frac{\Gamma}{2C(t;0)}, 0)$ follows similarly as above. We plug (ξ^*,η^*) into h and obtain Equation (46), finishing the proof. □

5 Solution of the optimization problem

In the previous sections we collected the building blocks for solving the Optimization Problem (OPT). The goal of this section is to prove that (OPT) is indeed solved by the candidate optimal control $u^* = (\xi^*,\eta^*)$ and that the value function is given by the candidate value function w. Given the control u^*, we denote the process controlled by u^* by $X^* := X^{u^*}$. We proceed as follows. In Section 5.1, we obtain bounds for w, u^* and X^*. Using these bounds, we show that u^* is an admissible control. We prove the verification theorem in Section 5.2. For both tasks it is essential that the w and u^* are given in closed form. Firstly, this enables us to verify that the required moment bounds for u^* hold. Secondly, the bounds for u^* turn into an upper bound for X^* via Gronwall's inequality; this is required in order to prove that the terminal constraint (Definition 2.1 (iv)) is fulfilled. Finally, the closed form solutions are key for the verification argument: Because of the conjectured singularity of the value function at terminal time T, we have to consider the limit $s \to T$ and exchange integration and limits; the main step in this respect is accomplished in Lemma 5.5.

5.1 Admissibility of the candidate optimal control

Using Equation (26), it can be shown by direct computation of the respective partial derivative that $C_1(t,\cdot)$ is strictly decreasing in $[t,T]$ (for details, see Kratz [2011] again, cf. Footnote 4). By the series expansion of coth, Equation (4) and Equation (30) we obtain the following inequalities. For $t \in [0,T)$ and $s_1, s_2 \in (t,T)$, $s_1 < s_2$,

$$0 < C(t;\theta) = C_1(t,T) < C_1(t,s_2) < C_1(t,s_1) < C_1(t,t) = C(t;0) \leq \frac{\Lambda}{T-t} + \sqrt{\alpha\sigma^2\Lambda}. \tag{47}$$

Using Taylor's theorem with integral remainder term, we can deduce the following useful bounds for the candidate value function w and the candidate optimal absolutely continuous control ξ^*.

Proposition 5.1. *Let $t \in [0,T)$ and $x \neq 0$. Then*

$$C_1(t,T)x^2 < w(t,x) \leq C(t;0)x^2 \leq \Big(\frac{\Lambda}{T-t} + \sqrt{\alpha\sigma^2\Lambda}\Big)x^2, \tag{48}$$

$$\frac{C_1(t,T)}{\Lambda}|x| < |\xi^*(t,x)| \leq \frac{C(t;0)}{\Lambda}|x| \leq \Big(\frac{1}{T-t} + \sqrt{\frac{\alpha\sigma^2}{\Lambda}}\Big)|x|. \tag{49}$$

Proof. Note first that $w(t,0) = \frac{\partial w}{\partial x}(t,0) = 0$ by Equations (22) and (34) (recall that $g(t,0) = t$). Thus,

$$w(t,x) = w(t,0) + \frac{\partial w}{\partial x}(t,0)x + \int_0^x \frac{\partial^2 w}{\partial x^2}(t,y)(x-y)dy = 2\int_0^x C_1(t,g(t,y))(x-y)dy$$

(cf. the proof of Theorem 4.7 for the second equality). Using (47), we can directly deduce (48). Furthermore, $\frac{\partial w}{\partial x}(t,x) = 2\Lambda\xi^*(t,x)$. Thus,

$$2\Lambda\xi^*(t,x) = \frac{\partial w}{\partial x}(t,0) + \int_0^x \frac{\partial^2 w}{\partial x^2}(t,y)dy = 2\int_0^x C_1(t,g(t,y))dy,$$

and (49) follows from (47). □

Using Gronwall's inequality, we can deduce an upper bound X^*.

Corollary 5.2. *Let $t \in [0,T)$ and $x \in \mathbb{R}$. Then*

$$|X^*(s)| \leq |x| \exp\Big(-\int_t^s \frac{C_1(u,T)}{\Lambda}du\Big) = |x| \exp\big(\frac{\Lambda\theta}{2}(s-t)\big)\frac{\sinh\big(\frac{\tilde\theta}{2}(T-s)\big)}{\sinh\big(\frac{\tilde\theta}{2}(T-t)\big)}.$$

Proof. We define the process $(\tilde{X}(s))_{s\in[t,T)}$ as the solution of the initial value problem

$$X'(s) = -\xi^*(s, X(s)), \quad X(t) = x.$$

By the structure of the process η^* (cf. Equation (45)), we have that for all $s \in [0,T)$, $|X^*(s)| \leq |\tilde{X}(s)|$. Therefore, by Gronwall's inequality, Proposition 5.1 and Equation (30),

$$|X^*(s)| \leq |\tilde{X}(s)| \leq |x| \exp\Big(- \int_t^s \frac{C_1(u,T)}{\Lambda} du\Big) = |x| \exp\big(\frac{\Lambda\theta}{2}(s-t)\big) \frac{\sinh\big(\frac{\tilde{\theta}}{2}(T-s)\big)}{\sinh\big(\frac{\tilde{\theta}}{2}(T-t)\big)}$$

as required. ☐

The bounds derived above allow us to deduce admissibility u^*.

Proposition 5.3. *Let $t \in [0,T)$, $x \in \mathbb{R}$ be the state at time t and $u^* = (\xi^*, \eta^*)$ be as in Equations* (44) *and* (45)*. Then $u^* \in \mathbb{A}(t,x)$.*

Proof. Note first that (i) of Definition 2.1 is straightforward and that (iii) holds as $\theta > 0$. By Proposition 5.1, Corollary 5.2 and Equation (4) we have

$$\begin{aligned} |\xi^*(s, X^*(s))| &\leq \frac{C(s;0)}{\Lambda} |X^*(s)| \leq |x| \sqrt{\frac{\alpha\sigma^2}{\Lambda}} \exp\big(\frac{\Lambda\theta}{2}(s-t)\big) \frac{\sinh\big(\frac{\tilde{\theta}}{2}(T-s)\big)}{\sinh\big(\frac{\tilde{\theta}}{2}(T-t)\big)} \frac{\cosh\big(\sqrt{\frac{\alpha\sigma^2}{\Lambda}}(T-s)\big)}{\sinh\big(\sqrt{\frac{\alpha\sigma^2}{\Lambda}}(T-s)\big)} \\ &\longrightarrow |x| \sqrt{\frac{\alpha\sigma^2}{\Lambda}} \exp\big(\frac{\Lambda\theta}{2}(T-t)\big) \frac{1}{\sinh\big(\frac{\tilde{\theta}}{2}(T-t)\big)} < \infty \quad \text{as } s \to T- . \end{aligned}$$

Thus, $\mathbb{E}[\int_t^T |\xi^*(s, X^*(s))|^4 ds] < \infty$. As $|\eta^*(t, X^*(s-))| < |X^*(s-)|$, we have $\mathbb{E}[\int_t^T |\eta^*(s, X^*(s-))|^8 ds] < \infty$ by Corollary 5.2. Finally, again by Corollary 5.2, $\lim_{s\to T-} X^*(s) = 0$. ☐

5.2 Verification

We are now finally able to prove that the Optimization Problem (OPT) is solved by the candidate optimal control in Equations (44) and (45) and that the value function is given as in Equation (22).

Theorem 5.4. *The value function of the Optimization Problem* (OPT) *is given by*

$$v(t,x) = C_1(t, g(t,x))x^2 + C_2(t, g(t,x))|x| + C_3(t, g(t,x))$$

for $t \in [0,T)$, $x \in \mathbb{R}^n$ (cf. Equation (22)*) where C_1, C_2, C_3 and g are given as in Section 4. The $\mathbb{P} \otimes \lambda$ - a.s. unique optimal control is given by $u^* = (\xi^*, \eta^*)$ as in Equations* (44) *and* (45)*, respectively.*

For the proof of the theorem, we require the following lemma, which relies on the fact that the value function v is an upper bound for the value function for $\Gamma = 0$ given in Equation (3).

Lemma 5.5. *Let $t \in [0,T)$, $x \in \mathbb{R}$ and $u \in \mathbb{A}(t,x)$. Then $\lim_{s\to T-} \mathbb{E}\big[w(s, X^u(s))\big] = 0$.*

Proof. Note first that there exists a constant $K > 0$ such that

$$\frac{K}{T-t} x^2 \leq C(t;\theta)x^2 \leq v(t,x), \tag{50}$$

where $C(t;\theta)$ is as in Equation (4). A control $u \in \mathbb{A}(t,x)$ has finite costs $J(t,x,u) < \infty$ by Definition 2.1 (ii) and Proposition 2.2. Let now $s \in (t,T)$. Then

$$\begin{aligned} J(t,x,u) &= \mathbb{E}\Big[\int_t^s \Big(\Lambda\xi(r)^2 + \theta\Gamma|\eta(r)| + \alpha\sigma^2 X^u(r)^2\Big) dr\Big] + \mathbb{E}\Big[\int_s^T \Big(\Lambda\xi(r)^2 + \theta\Gamma|\eta(r)| + \alpha\sigma^2 X^u(r)^2\Big) dr\Big] \\ &\geq \mathbb{E}\Big[\int_t^s \Big(\Lambda\xi(r)^2 + \theta\Gamma|\eta(r)| + \alpha\sigma^2 X^u(r)^2\Big) dr + v(s, X^u(s))\Big] \\ &\overset{(50)}{\geq} \mathbb{E}\Big[\int_t^s \Big(\Lambda\xi(r)^2 + \theta\Gamma|\eta(r)| + \alpha\sigma^2 X^u(r)^2\Big) dr\Big] + K\mathbb{E}\Big[\frac{X^u(s)^2}{T-s}\Big]. \end{aligned}$$

Thus by the monotone convergence theorem,

$$\begin{aligned}
J(t,x,u) &\geq \lim_{s\to T-}\Big(\mathbb{E}\Big[\int_t^s\Big(\Lambda\xi(r)^2+\theta\Gamma|\eta(r)|+\alpha\sigma^2X^u(r)^2\Big)dr\Big]+K\mathbb{E}\Big[\frac{X^u(s)^2}{T-s}\Big]\Big)\\
&= J(t,x,u)+K\lim_{s\to T-}\mathbb{E}\Big[\frac{X^u(s)^2}{T-s}\Big];
\end{aligned}$$

hence $\lim_{s\to T-}\mathbb{E}\big[\frac{X^u(s)^2}{T-s}\big]=0$. The assertion now follows directly from Proposition 5.1 (cf. Inequality (48)). ☐

Proof of Theorem 5.4. Let $t\in[0,T)$, $x\in\mathbb{R}$, $u\in\mathbb{A}(t,x)$ and $s\in[t,T)$. We apply Itô's formula to the function $w(r,X^u(r))$ (recall that w is differentiable by Theorem 4.4):

$$\begin{aligned}
w(t,x) &= w(s,X^u(s))+\int_t^s\Big(\frac{\partial w}{\partial x}(r,X^u(r))\xi(r)-\frac{\partial w}{\partial r}(r,X^u(r))\Big)dr\\
&\qquad+\int_t^s\big(w(r,X^u(r-))-w(r,X^u(r-)-\eta(r))\big)\pi(dr)\\
&\leq w(s,X^u(s))+\int_t^s f(\xi(r),\eta(r),X^u(r))dr+\int_t^s\big(w(r,X^u(r-))-w(r,X^u(r-)-\eta(r))\big)\pi(dr)\\
&\qquad-\int_t^s\theta\big(w(r,X^u(r-))-w(r,X^u(r-)-\eta(r))\big)dr \qquad (51)\\
&= w(s,X^u(s))+\int_t^s f(\xi(r),\eta(r),X^u(r))dr+\int_t^s\big(w(r,X^u(r-))-w(l,r,X^u(r-)-\eta(r))\big)M(dr), \qquad (52)
\end{aligned}$$

where M is the compensated Poisson process given by $M(r):=\pi(r)-\theta r$ and Inequality (51) follows from Theorem 4.8, with equality if and only if $\lambda[u=u^*]=0$. Taking expectations on both sides, we obtain

$$\begin{aligned}
w(t,x) &\leq \mathbb{E}[w(s,X^u(s))]+\mathbb{E}\Big[\int_t^s f(\xi(r),\eta(r),X^u(r))dr\Big]\\
&\qquad+\mathbb{E}\Big[\int_t^s w(r,X^u(r-))-w(r,X^u(r-)-\eta(r))M(dr)\Big], \qquad (53)
\end{aligned}$$

with equality if and only if $\lambda\otimes\mathbb{P}[u=u^*]=0$. By Proposition 5.1, there exists a constant $K=K(s)$ such that

$$\mathbb{E}\Big[\int_t^s|w(r,X^u(r-))-w\big(r,X^u(r-)-\eta(r,X^u(r-))\big)|^2dr\Big]\leq K(s)\Big(\mathbb{E}\Big[\int_t^s\big(|X^u(r-)|^4+|\eta(r)|^4\big)dr\Big]\Big)<\infty$$

by Definition 2.1 (ii) and Proposition 2.2. As $\langle M\rangle(s)=\theta s$, this implies that the stochastic integral in Equation (52) is a martingale. Thus, taking the limit $s\to T-$ in Inequality (53), we obtain by Lemma 5.5 and the monotone convergence theorem that $w(t,x)\leq\mathbb{E}[\int_t^T f(\xi(r),\eta(r),X^u(r))dr]=J(t,x,u)$, again with equality if $\lambda\otimes\mathbb{P}[u=u^*]=0$.

For uniqueness, let $u=(\xi,\eta),\tilde{u}=(\tilde{\xi},\tilde{\eta})\in\mathbb{A}(t,x)$ and $\mu\in(0,1)$. We define the convex combination $\bar{u}=(\bar{\xi},\bar{\eta})$:

$$\bar{\xi}(s)=\mu\xi(s)+(1-\mu)\tilde{\xi}(s),\quad \bar{\eta}(s)=\mu\eta(s)+(1-\mu)\tilde{\eta}(s)$$

for $s\in[t,T)$. Thus, $X^{\bar{u}}(s)=\mu X^u(s)+(1-\mu)X^{\tilde{u}}(s)$ and $\bar{u}\in\mathbb{A}(t,x)$. Notice that

$$\mathbb{P}\otimes\lambda\big[u\neq\tilde{u}\big]>0\quad\text{implies}\quad\mathbb{P}\otimes\lambda\big[\xi\neq\tilde{\xi}\big]>0 \qquad (54)$$

as else $\mathbb{P}[\lim_{s\to T-}X^u(s)\neq\lim_{s\to T-}X^{\tilde{u}}(s)]>0$, a contradiction to Definition 2.1 (iv). Hence by the convexity of f,

$$\begin{aligned}
J(t,x,\bar{u}) &= \mathbb{E}\Big[\int_t^T f\big(\bar{\xi}(r),\bar{\eta}(r),X^{\bar{u}}(r)\big)dr\Big]\\
&\leq \mathbb{E}\Big[\int_t^T \mu f\big(\xi(r),\eta(r),X^u(r)\big)+(1-\mu)f\big(\tilde{\xi}(r),\tilde{\eta}(r),X^{\tilde{u}}(r)\big)dr\Big]=\mu J(t,x,u)+(1-\mu)J(t,x,\tilde{u});
\end{aligned}$$

by the strict convexity of f in the first argument and (54), we have equality if and only if $u=\tilde{u}$ $\mathbb{P}\otimes\lambda$ - a.s. ☐

6 Properties and application

In this section, we discuss the properties of the value function v and the optimal control u^*. In Section 6.1, we discuss the dependence of u^* and v on Γ; this is enabled by the fact that both are given in closed form. In Section 6.2, we analyze the case $\alpha\sigma^2 = 0$ which we had excluded in Sections 4 and 5. We conclude by motivating the optimization problem by applying it to a financial market model in Section 6.3.

6.1 $\alpha\sigma^2 > 0$

Theorem 5.4 confirms the structure of the optimal control and the value function of the Optimization Problem (OPT) we have expected in the heuristics of Section 3 (cf. also Figure 1).

Both for large initial states x ($|x| \geq \bar{X}(t,T)$) and for small initial states ($|x| \leq \bar{X}(t,t) = \beta(t) = \frac{\Gamma}{2C(t;0)}$), v is a quadratic polynomial. In between, it is an "interpolation" of these polynomials. The value function and the optimal control for $|x| \leq \frac{\Gamma}{2C(t;0)}$ are the same as the ones for $\theta = 0$ (in particular, $\eta^* = 0$). For larger states, the absolute value of η^* is positive; after the jump of the Poisson π process at time s, the controlled process is $X^*(s) = \frac{\Gamma}{2C(s;0)}$. The optimal trajectory until a jump of π for an initial state $|x| = \bar{X}(t,s) \in (\bar{X}(t,t), \bar{X}(t,T))$ is given by the function $\bar{X}(\cdot, s)$ in $[t,s]$.

It is interesting to examine the dependence of v and u^* on Γ. Costs are higher for larger Γ and therefore the value function increases in Γ (cf. the Cost Functional (2)). Furthermore, the jump control η is less attractive for large Γ and therefore η^* should decrease in Γ whereas the optimal absolutely continuous control ξ^* should increase in Γ. The following proposition confirms these intuitions. Additionally, (i) of the proposition shows that the costs for the jump control, $\theta\Gamma|\eta^*|$, are negligible for large initial states $|x| >> \bar{X}(t,T)$ as in this case $v(t,x) \approx C_1(t,T)x^2$ which is independent of Γ. To stress the dependence on Γ, we add it as an argument for the remainder of the section and write $u^*(t,x,\Gamma) := u^*(t,x)$, etc.

Proposition 6.1. *Let $t \in [0,T)$, $s \in [t,T]$ and $x \in \mathbb{R}$. Then*

(i) $C_1(t,s,\cdot)$ is constant, $C_2(t,s,\cdot)$ is strictly increasing and $C_3(t,s,\cdot)$ is strictly decreasing.

(ii) $\bar{X}(t,s,\cdot)$ is strictly increasing.

(iii) $|\xi^*(t,x,\cdot)|$ is strictly increasing and $|\eta^*(t,x,\cdot)|$ is strictly decreasing on the interval $(0, 2|x|C(t;0))$ and constant on the interval $(2|x|C(t;0), \infty)$.

Proof. (i), (ii) The assertions follow directly from Proposition 4.1 respectively Equations (30) - (33).

(iii) Note first that $|\eta^*(t,x,\Gamma)| > 0$ if and only if $\Gamma \in (0, 2|x|C(t;0))$ (cf. Equation (45)). Hence both controls are constant for $\Gamma > 2|x|C(t;0)$.

For monotonicity of ξ^*, let without loss of generality $x > 0$ and $\Gamma < \tilde{\Gamma} < 2xC(t;0)$. By (ii) (cf. Equation (24)), $g(t,y,\tilde{\Gamma}) \leq g(t,y,\Gamma)$ for all $y \in [0,x]$ with strict inequality for $\frac{\tilde{\Gamma}}{2C(t;0)} < y < \bar{X}(t,T,\Gamma)$. Therefore (cf. the proof of Proposition 5.1),

$$\xi^*(t,x,\Gamma) = \frac{1}{\Lambda}\int_0^x C_1(t, g(t,y,\Gamma), \Gamma)dy < \frac{1}{\Lambda}\int_0^x C_1(t, g(t,y,\tilde{\Gamma}), \tilde{\Gamma})dy = \xi^*(t,x,\tilde{\Gamma})$$

by (i) and the fact that C_1 is strictly decreasing in s (cf. the discussion at the beginning of Section 5.1).

Monotonicity of $|\eta^*|$ follows directly from Equation (45).

□

In Figure 2, we compare the optimal control for $\Gamma > 0$ (left picture) and $\Gamma = 0$ (right picture) for a numerical example. In both pictures, thick solid lines denote realized trajectories of X^*. For those scenarios where π jumps at time $\tau < T$, the dotted lines refer to the continuation of the trajectory without jump. The dashed line in the left picture corresponds to the boundary β and the thin solid line in the right picture corresponds to the case $\theta = 0$. For the larger initial state, the boundary β is never crossed unless π jumps (upper solid line respectively dotted lines in the left picture). At all times, the optimal jump control is such that after a jump of π, the state is exactly on the boundary. Afterwards, the controlled process X^* stays (strictly) below the boundary for the entire horizon. In comparison to the optimal control for $\Gamma = 0$ (thick solid line in the right picture), the absolutely continuous control is larger (cf. Proposition 6.1 (iv)); it is however smaller than the control for



Figure 2: Optimal controls for $\Gamma > 0$ (left picture) and $\Gamma = 0$ (right picture). In both pictures, thick solid lines denote realized trajectories of X^*. For those scenarios where the Poisson process jumps at time $\tau = 0.4 < T = 1$, the dotted lines refer to the continuation of the trajectories without jumps. The dashed line in the left picture corresponds to the boundary β and the thin solid line in the right picture corresponds to optimal control for $\theta = 0$. The larger initial state is $x = 1.2$; the smaller one is $x = 0.3$ (which results in $s = 0.1$). Furthermore, $\Lambda = 2.5$, $\alpha = 4$ $\sigma = 1$, $\theta = 3$ $\Gamma = 2$.

$\theta = 0$. The introduction of the jump control decreases ξ^* (see Kratz and Schöneborn [2012b]); this effect is decreased significantly by the introduction of the absolute value costs for the jump control. Furthermore, the optimal jump control is smaller than for $\Gamma = 0$, where $\eta(t, x) = x$ is optimal. For the smaller initial state, the boundary β is crossed at time s, i.e., the optimal trajectory $X^*(t)$ until s is $\bar{X}(t, s)$ (unless the Poisson process jumps before). After the controlled process crosses the boundary it stays (strictly) below it until the end of the horizon.

6.2 $\alpha\sigma^2 = 0$

In Sections 4 and 5 we excluded the case $\alpha\sigma^2 = 0$. The main reason was that in some formulae the term $\alpha\sigma^2$ appears in the denominator (cf., e.g., Equations (29) and (33)) and it would thus have complicated the exposition of the results. However, the case $\alpha\sigma^2 = 0$ can be treated in a similar way as the case $\alpha\sigma^2 > 0$.

We define C_1, C_2, C_3 and $\bar{X}$ as before by the Initial Value Problems (18) - (21). Note that only the differential equation for C_1 depends on $\alpha\sigma^2$. Therefore C_2, C_3 and $\bar{X}$ only depend on $\alpha\sigma^2$ through C_1. We can compute solutions of the initial value problems directly or by taking the limits for $\alpha\sigma^2 \to 0$ in Equations (26) - (29).

Note first that $\alpha\sigma^2 = 0$ implies $\tilde{\theta} = \theta$ and $C(t; 0) = \frac{\Lambda}{T-t}$. Therefore,

$$\bar{X}(t, t) = \beta(t) = \frac{\Gamma}{2C(t; 0)} = \frac{\Gamma}{2\Lambda}(T - t).$$

We can compute the partial derivative of $\bar{X}$ with respect to s and obtain $\frac{\partial \bar{X}}{\partial s}(t, s) = 0$; therefore,

$$\bar{X}(t, s) = \bar{X}(t, t) = \frac{\Gamma}{2\Lambda}(T - t) \tag{55}$$

for $t \in [0, T)$, $s \in [t, T]$. Hence, we expect only two different control regions. We expect that $|\eta^*| > 0$ only for $|x| > \frac{\Gamma}{2\Lambda}(T - t)$ and that the value function and the optimal control are given by

$$v(t, x) = \begin{cases} C(t; 0)x^2 = \frac{\Lambda}{T-t}x^2 & \text{if } |x| \leq \frac{\Gamma}{2\Lambda}(T - t) \\ C_1(t, T)x^2 + C_2(t, T)|x| + C_3(t, T) & \text{if } |x| > \frac{\Gamma}{2\Lambda}(T - t) \end{cases} \tag{56}$$

respectively

$$\xi^* = \xi^*(t, x) := \frac{1}{2\Lambda}\frac{\partial w}{\partial x}(t, x) = \begin{cases} \frac{2C_1(t,T)x + \operatorname{sgn}(x)C_2(t,T)}{2\Lambda} & \text{if } |x| > \frac{\Gamma}{2\Lambda}(T - t) \\ \frac{x}{T-t} & \text{if } |x| \leq \frac{\Gamma}{2\Lambda}(T - t), \end{cases} \tag{57}$$

$$\eta^* = \eta^*(t, x) := \begin{cases} \operatorname{sgn}(x)\Big(|x| - \frac{\Gamma}{2\Lambda}(T - t)\Big) & \text{if } |x| > \frac{\Gamma}{2\Lambda}(T - t) \\ 0 & \text{if } |x| \leq \frac{\Gamma}{2\Lambda}(T - t), \end{cases} \tag{58}$$

All steps of the proof of Theorem 5.4 can be replicated for the case $\alpha\sigma^2 = 0$ in a straightforward manner. The solution of the Optimization Problem (OPT) is given by Equations (57) and (58); the value function is given as in Equation (56).



Figure 3: The left and the middle picture illustrate the same scenarios as Figure 2 for $\alpha\sigma^2 = 0$. In addition, an intermediate initial state is displayed ($x = 0.6$). The right picture illustrates the fact that $\xi^*(t,\cdot)$ is not differentiable at $x = \frac{\Gamma T}{2\Lambda}$. The parameters are the same as in Figure 2.

We illustrate the structure of the optimal control in the left and the middle picture of Figure 3. All lines have the same meaning as in Figure 2; we only add a realized trajectory for an intermediate initial state which is only slightly larger than the boundary $\beta(0) = \frac{\Gamma}{2\Lambda}T$. We observe that the optimal trajectory $X^*(s)$ never crosses the boundary $\frac{\Gamma}{2\Lambda}(T-s)$ if π does not jump. Let us briefly comment on this structure. For $\alpha\sigma^2 = \theta = 0$, the optimal absolutely continuous control is constant (see, e.g., Kratz and Schöneborn [2012b]; cf. also the thin solid line in the middle picture of Figure 3 which refers to $\alpha\sigma^2 = \theta = 0$). The boundary itself is linear and it is the trajectory of the optimal control (for $\theta = 0$) with initial state *on* the boundary. If the initial state is above the boundary, $\theta > 0$ *decreases* the optimal absolutely continuous control, i.e., if π does not jump, the trajectory is concave and thus never crosses the boundary. If π jumps (in the displayed scenario at time τ), the resulting state is again on the boundary. However, it does *not cross* it but stays on it until the end of the horizon. Finally, the boundary β is decreasing in $\alpha\sigma^2$ (cf., e.g., Equation (13)). This results in the small state in the left picture to be below the boundary from the beginning (cf. Figure 2, where the same initial state was above the respective boundary). Again, the optimal absolutely continuous control is constant in this case.

By analyzing Equation (57), we can observe that $\xi^*(t,\cdot)$ is continuous but *not* differentiable at $|x| = \beta(t) = \frac{\Gamma}{2\Lambda}(T-t)$. The right picture of Figure 3 illustrates this property.

6.3 Optimal liquidation in dark pools with adverse selection

The Optimization problem (OPT) is motivated by a cost minimizing problem in finance. In the following, we describe this application.

In recent years, equity trading has been transformed by the advent of so-called dark pools. These alternative trading venues differ significantly from classical exchanges and have gained a considerable market share, especially in the US. They generally share the following two stylized facts. First, the liquidity available in dark pools is not quoted, hence making trade execution uncertain and unpredictable. Second, dark pools do not determine prices. Instead, they monitor the prices determined by the classical exchanges and settle trades in the dark pool only if possible at these prices (see Mittal [2008] and Degryse et al. [2009] for further details and a typology of dark pools). In traditional exchanges, large transactions can result in adverse price movements due to a lack of liquidity in the market or because of the information revealed by them. Trades in the dark pool have no or less price impact. On the other hand, order execution in dark pools is uncertain and cannot be enforced. Additionally, dark pool orders incorporate indirect costs by so-called *adverse selection*: Liquidity seeking traders find that their trades in the dark pool are usually executed just before a favorable price move in the market, i.e., exactly when they do not want them to be executed since they miss out on the price improvement (see Mittal [2008] for explanations for adverse selection; Hendershott and Mendelson [2000] find theoretical evidence that adverse selection arises due to information asymmetries in the market).

In the following, we consider a risk averse investor who has to liquidate a position $x \in \mathbb{R}$ in a risky asset within a finite time horizon $[0,T]$. The investor has the opportunity to trade at a traditional exchange by controlling her trading intensity ξ; due to linear price impact (see Almgren and Chriss [2001]) this yields quadratic price impact costs $\int_0^T \Lambda\xi(s)ds$. Additionally she can place orders η in a dark pool; we model order execution by the Poisson process π: the orders are executed fully at the jump times of π. While no price impact costs have to be paid for dark pool orders, indirect costs $\int_0^T \theta\Gamma|\eta(s)|ds$ arise from adverse selection. Kratz and Schöneborn [2012a] justify this cost term for adverse selection for the discrete time analogue of the market model; for the continuous time model this is accomplished by Naujokat and Horst [2011]. We can interpret the controlled process X^u ($u = (\xi,\eta)$) as the stock holdings of the investor; the liquidation constraint is reflected by Definition 2.1 (iv).

The last term in the additive penalty function, $\alpha \int_t^T \sigma^2 X^u(s)^2 ds$, reflects the market risk of the position and penalizes slow liquidation; here, $\alpha \geq 0$ is the personal risk aversion of the investor and σ^2 is the variance of the price process of the asset (this expression for the market risk is again due to Almgren and Chriss [2001]). The case $\alpha\sigma^2 = 0$ treated in Section 6.2 hence describes the important special case of a risk neutral investor.

The Optimization Problem (OPT) hence reflects the trade-off of price impact costs versus execution uncertainty and adverse selection costs faced by the investor. As discussed in Sections 6.1 and 6.2, adverse selection changes the optimal strategy in two main respects: Instead of placing the full remainder of the position in the dark pool (as it is the case for optimal liquidation without adverse selection, see Kratz and Schöneborn [2012b]; cf. also the right picture of Figure 2) a smaller fraction is offered in the dark pool as the investor does not want to miss out on the expected favorable impending price move completely; for small positions, the dark pool is not used at all. On the other hand, the trading intensity in the traditional exchange is increased.

References

R. Almgren and N. Chriss. Optimal execution of portfolio transactions. *Journal of Risk*, 3(2):5–39, 2001.

F. E. Benth and K. Reikvam. A connection between singular stochastic control and optimal stopping. *Applied Mathematics and Optimization*, 49(1):27–41, 2004.

H. Degryse, M. Van Achter, and G. Wuyts. Shedding light on dark liquidity pools. *The Institutional Investor*, 2009(1):147–155, 2009.

F. B. Hanson. *Applied stochastic processes and control for jump-diffusions: modeling, analysis, and computation.* SIAM, Philadelphia, 2007.

T. Hendershott and H. Mendelson. Crossing networks and dealer markets: Competition and performance. *Journal of Finance*, 55(5):2071–2115, 2000.

P. Kratz. *Optimal liquidation in dark pools in discrete and continuous time.* PhD thesis, Humboldt University Berlin, 2011.

P. Kratz and T. Schöneborn. Optimal liquidation in dark pools. *Preprint*, 2012a.

P. Kratz and T. Schöneborn. Portfolio liquidation in dark pools in continuous time. *Preprint*, 2012b.

H. Mittal. Are you playing in a toxic dark pool? A guide to preventing information leakage. *Journal of Trading*, pages 20–33, 2008.

F. Naujokat and U. Horst. When to cross the spread: Curve following with singular control. *Preprint*, 2011.

F. Naujokat and N. Westray. Curve following in illiquid markets. *Mathematics and Financial Economics*, 4(4): 299–335, 2011.

B. Øksendal and A. Sulem. *Applied stochastic control of jump diffusions.* Springer, Berlin, 2007.

H. Pham. *Continuous-time stochastic control and optimization with financial applications.* Springer, Berlin, 2009.

A. Schied, T. Schöneborn, and M. Tehranchi. Optimal basket liquidation for CARA investors is deterministic. *Applied Mathematical Finance*, 17(6):471–489, 2010.